

Post Operative Instructions Capsular Release Surgery

In general, rehabilitation after arthroscopic shoulder surgery can be divided into three phases:

Phase 1 consists of the first few weeks after surgery and involves decreasing swelling and inflammation after the operation. Immediately after the surgery you will ice the shoulder as much as possible. At your first office visit, I will remind you of your “5 for 5” exercise program.

Phase 2 consists of regaining motion. Often after a capsular release you will have very near normal motion; your goal is to keep what we achieved in surgery.

Phase 3 consists of getting the strength back in the arm and getting the shoulder “back to normal.” This time period continues until about 1 full year after your surgery. We may have you participate in a formal physical therapy program during this time.

What to Expect Immediately after Surgery:

- Swelling:** Ice the shoulder for 20 minutes every two hours for the first 2-3 days after surgery. If you received an ‘ice machine,’ use this as much as possible.
- Bandage/Immobilizer:** The bandages can be removed the day after surgery and replaced with band-aids. Stop using your sling as soon as your arm ‘wakes up’ from the nerve block (if applicable).
- Exercises/Range of Motion:** At your first post-op visit you will be taught a few exercises and stretches (“5 for 5”); the goal is to maintain motion achieved in surgery.
- Medications:** You will be given an anti-inflammatory medicine to prevent recurrence of scar tissue; usually this is taken for 6 weeks. Do not take any additional Tylenol/acetaminophen if your prescribed pain medication contains Tylenol (acetaminophen or APAP). Take your pain medication as directed. Narcotics can cause constipation; you may need to use a laxative and/or stool softener.
- Eating/Showering:** Your first meal should be clear liquids only (water, tea, broth). If you tolerate that, progress to regular food at your next meal. You may shower if you can keep the wounds dry: place a waterproof bandage over the wound and face shower so that the water does not hit your arm. If you can’t keep the wound dry, then wait until the staples/stitches have been removed.
- Driving/Returning to Work:** You may drive when you are off narcotic pain pills and feel safe driving. Use your judgment and be safe. Return to work depends on how you feel. If you work at a desk job or in a supervisory role, you may return as early as 1 week after surgery. If you do heavy lifting or overhead activity you may need more time off. If there was no repair, you can use the arm as tolerated, even if it is sore.
- Caring For Incision:** Keep your incision clean and dry. Do not apply any lotions to the wound.
- Follow-up Appointment:** _____
If there is no appointment time and date written on the line above, please call office (804) 288-3136 and ask for The Walker Team to make an appointment 10-12 days post op.
- Therapy:** Need for therapy will be determined on a case-by-case basis. Protocols are given to home therapists and to outpatient therapists based on your operation.

Symptoms to Report:

Notify Dr. Walker's office immediately if you develop any of the following:

- Fever above 101.5 degrees Fahrenheit
- Nausea or vomiting
- Inability to urinate or urinary incontinence
- Changes in sensations in your arms/legs
- Pain not relieved by medication
- Redness and or excessive swelling
- Drainage from incision

We are here to help. Feel free to contact our office anytime to ask any questions or concerns. Call (804) 288-3136 and ask for "The Walker Team." You may also reach us by email through the Patient Portal System (Oberd) or by going on to our website www.virginiashoulder.com.